

An on-going multi-wavelength survey of Ly α emitters at redshifts $z = 2 - 8$

Kim K. Nilsson,^{1,2} Johan P.U. Fynbo², Palle Møller¹ and Alvaro Orsi³

¹*ESO Garching, Karl-Schwarzschild-Strasse 2, D-85748 Garching bei München, Germany*

²*Dark Cosmology Centre, Juliane Maries Vej 30, DK-2100 København Ø, Denmark*

³*Department of Astronomy, Pontificia Universidad Católica, Casilla 306, Santiago 22, Chile*

Abstract. In the last decade, the technique of finding Ly α -emitters through narrow-band imaging has become a promising method of detecting high redshift galaxies. Ly α -emitters have been found from redshifts $z \gtrsim 2$, up to the highest redshift source known to date at $z = 6.96$. Several surveys are also underway to find $z = 7 - 9$ sources. But these very high redshift sources are too faint to be studied in great detail, and more information can be found from studying the same class of objects at lower redshifts. Here we present our survey strategy to determine the nature of Ly α -emitters at lower redshifts, through multi-wavelength surveys, and our plans to extend the survey to redshift $z = 8.8$.

1. Introduction

The success of narrow-band searches, focusing on Ly α , was predicted by Partridge & Peebles (1967) nearly 40 years ago, but early surveys failed to produce anything other than upper limits. The unexpected faintness of the objects caused it to take almost three decades before the narrow-band technique was successfully used to identify the first high redshift Ly α emitting galaxies (Møller & Warren 1993; Hu & McMahon 1996), and it is only recently, with the advent of 8 m class telescopes and sensitive detectors, that larger samples of Ly α selected objects have been reported (Steidel et al. 2000; Malhotra & Rhoads 2002; Fynbo et al. 2003; Venemans et al. 2005). Lately, much of the attention has been turned to detecting Ly α -emitters in the redshift range of $z = 4 - 7$, and the lower redshifts have been neglected. Due to the increase in luminosity distance, objects at these redshifts are very difficult to study in greater detail, once they are found. At lower redshifts (i.e. $z = 2 - 3$), it is still possible to detect, or at least put strong upper limits on, the SED of the galaxy. Hence at these redshifts, we can study the nature of Ly α selected galaxies, using public multi-wavelength surveys. However, Ly α also appears to be an excellent tool in finding very high redshift sources (e.g. Iye et al. 2006) and so it is also of interest to push the limits of what is observable. We can then use the strength of wide-field narrow-band imaging for Ly α to find very high redshift sources

($z > 8$), which in turn will enable a study of when re-ionisation happened and how galaxies formed.

In these proceedings, we present our on-going survey strategy to study the nature of Ly α -emitters at lower redshifts, and present a future Public Survey that we expect will find several very high redshift sources. Magnitudes are in the AB system.

Survey plans

In order to understand the physical nature of any high redshift galaxy, multi-wavelength coverage enabling SED fitting analysis is crucial. Hence recent large surveys, such as the GOODS fields (Giavalisco et al. 2004) and the COSMOS¹ survey are invaluable. These are the fields we target in our narrow-band surveys. We have nearly completed a survey at redshift $z = 3.15$ in the GOODS-S field (Nilsson et al. 2006; Nilsson et al. 2007, to be submitted). The narrow-band imaging we have in this field comes from the VLT/FORS instrument, with a field-of-view of $7' \times 7'$. In this field we have found 23 candidate compact Ly α -emitting candidates, and one so-called Ly α -blob. We have also started a similar survey in the GOODS-N field, using the Nordic Optical Telescope (NOT) on La Palma, Canary Islands, Spain. This survey is focused on two redshifts, $z = 2.34$ and 3.15 . The data shows 10 candidates in the lower redshift filter. The higher redshift filter imaging is not complete. Finally, we are proposing a Public Survey with the VISTA telescope; ELVIS - Emission Line galaxies with VISTA. This survey will be conducted with a narrow-band filter centred on Ly α at $z = 8.8$, see section 3.

2. GOODS-S and the Ly α -blob

In December 2002, one $7' \times 7'$ field, centred on RA $03^{\text{h}}32^{\text{m}}21.8^{\text{s}}$ and Dec $-27^{\circ}45'52''$ (J2000) was observed with VLT/FORS1. The field was observed for 8.3 hours in a narrow-band filter with central wavelength $5055 \text{ \AA}$ and with a FWHM of $60 \text{ \AA}$. This corresponds to a redshift range of $z = 3.126 - 3.174$ for Ly α . The images were reduced and calibrated with standard methods (see Nilsson et al. 2007, to be submitted, for details). The 5σ limiting magnitude for the narrow-band image is 25.8.

We selected objects using the HST/ACS F435W and F606W images available from GOODS and using SExtractor. The narrow-band image was used as a detection image, and broad-band magnitudes were measured in circular apertures placed at the position of the narrow-band sources on rebinned HST images. Objects with a measured equivalent width larger than zero and “signal-to-noise” of the equivalent width measurement larger than three were selected. Our final catalog includes 26 objects, of which three are associated with the Ly α -blob (sec. 2.1.; Nilsson et al. 2006) and the other 23 are compact Ly α -emitting candidates which will be presented in a forthcoming paper (Nilsson et al. 2007, to be submitted).

¹<http://www.astro.caltech.edu/~cosmos/>

2.1. The $\text{Ly}\alpha$ blob

In the narrow-band image, we found a so-called $\text{Ly}\alpha$ -blob, similar to what has been found by many authors before (e.g. Steidel et al. 2000; Matsuda et al. 2004). The blob has a size of approximately 70 kpc in projected diameter and a total $\text{Ly}\alpha$ line luminosity of $1.0 \cdot 10^{43} \text{ erg s}^{-1}$. A contour-plot, as well as a plot of the blob surface brightness and the spectrum can be found in figure 1. The total multi-wavelength data on the blob can be seen in figure 2.




Figure 1. *Left a)* Flux calibrated spectrum of the blob emission line. The line has the characteristic blue side absorption, indicating high redshift. *b)* The part of the spectrum where $\text{H}\beta$ and $[\text{OIII}]$ should have been observed if the emission line was $[\text{OII}]$ at a redshift of $z \approx 0.36$. These lines are not observed and therefore we conclude the observed line is due to $\text{Ly}\alpha$ at $z = 3.16$. *Middle* Contour-plot of narrow-band emission from the $\text{Ly}\alpha$ overlaid the HST V-band image. The narrow-band image has been continuum subtracted by subtracting the re-binned, smoothed and scaled HST/V-band image. Contour levels are $2 \cdot 10^{-4}$, $4 \cdot 10^{-4}$ and $6 \cdot 10^{-4} \text{ erg s}^{-1} \text{ cm}^{-2}$ in restframe flux (corresponding to $1.2 \cdot 10^{-18}$, $2.5 \cdot 10^{-18}$ and $3.7 \cdot 10^{-18}$ in observed flux). The image is $18'' \times 18''$ ($18''$ corresponds to a physical size of $\sim 133 \text{ kpc}$ at $z = 3.15$). The dotted lines indicate the slitlet position for our follow-up spectroscopy. *Right* Plot of surface brightness as function of radius. The flux is the sky subtracted narrow-band flux. The PSF of the image is illustrated by the solid line, and the dotted line is the best fit model of Dijkstra et al. (2006a,b). The deficit at $\sim 45 \text{ kpc}$ is due to the asymmetric appearance of the blob.




Figure 2. Thumbnail images of all available multi-wavelength data in the GOODS South field, centred on the $\text{Ly}\alpha$ blob. All images are $18'' \times 18''$.

There have been three proposed energy sources of blobs. These are *i)* hidden QSOs (Haiman & Rees 2001; Weidinger et al. 2005), *ii)* star formation

and superwinds from (possibly obscured) starburst galaxies (Taniguchi et al. 2001; Ohya et al. 2003), and *iii*) so-called cold accretion (Haiman, Spaans & Quataert 2000; Fardal et al. 2001; Sommer-Larsen 2005; Dijkstra et al. 2006a,b; Dekel & Birnboim 2006). Cooling flows are phenomena observed in galaxy clusters for more than a decade. These are explained by gas which is cooling much faster than the Hubble time through X-ray emission in the centres of the clusters. However, cooling emission from a galaxy, or a group sized halo can be dominated by Ly α emission, so-called cold accretion. Until the discovery of our GOODS-S blob, cold accretion had not been observed. Because of the extensive multi-wavelength coverage of our blob, we were able to conclude that it is powered by cold accretion.

The argument that the GOODS-S blob is powered by cold accretion comes from ruling out the other two options. As seen in the thumb-nail images in figure 2, there are no detected continuum counterparts to the blob Ly α emission. A starburst galaxy would have been observed in the deep optical or IR data we have, and an AGN would have been detected in the X-rays or the IR. Even an obscured AGN should be observed with the *Spitzer*/*MIPS* bands. There were two objects of particular interest to us; object #3 and object #6 in the contour-plot of figure 1. Object #3 since it appeared to have some faint Ly α -emission surrounding it, and had a photometric redshift similar to the blob ($z_{phot} = 2.9^{+1.41}_{-0.59}$), and object #6 since it only appeared in the *Spitzer* bands. Object #3 does appear to have some narrow-band flux with very low significance ($< 2\sigma$) and so we studied the UV properties of this galaxy. However, the measured UV flux of this galaxy is not sufficient to photo-ionise the blob. We considered that object #6 was a hidden AGN, but its IR colours are more indicative of a higher redshift starburst galaxy. After ruling out the possibility that the blob was powered by star formation, or AGN activity, we hence concluded that this blob must be the first observation of cold accretion of neutral gas onto a dark matter halo. For a more extensive discussion of this blob, see Nilsson et al. (2006).

3. ELVIS - Emission Line galaxies with VISTA Survey

Ly α -emitters is one of the most promising methods of finding very high redshift ($z > 6$) galaxies. To date, the galaxy with highest redshift known ($z = 6.96$, Iye et al. 2006) was found as a Ly α -emitter through narrow-band imaging techniques. Hence, there are now many proposed projects to find $z > 7$ Ly α -emitters. One such project is a Public Survey with the VISTA telescope. VISTA² is a new telescope being built near Paranal in Chile. It is a 4-m telescope, with a wide-field IR camera array. It is equipped with the standard set of broad-band filters, as well as a custom-made narrow-band filter with central wavelength 1185 nm and FWHM ≈ 10 nm purchased by the Dark Cosmology Centre (DK). VISTA will devote 85 % of its time for Public Surveys and will be run by ESO. The Public Survey we propose is called ELVIS – Emission Line galaxies with VISTA Survey (Nilsson et al. 2007, in prep.), now part of a larger survey called

²www.vista.ac.uk

Ultra-VISTA, which proposes to do both very deep broad- and narrow-band imaging.

The ELVIS part of the proposal, the deep narrow-band imaging, is planned to go to a depth of $3.7 \cdot 10^{-18}$ erg cm $^{-2}$ s $^{-1}$ (5σ) in an 0.9 square degree field. The science case is to find a large sample of emission-line galaxies at several redshifts. The observations will be sensitive to H α at redshift $z = 0.8$, [OIII] and H β at $z = 1.4$, [OII] at $z = 2.2$ and Ly α at redshift $z = 8.8$. For the Ly α -emitters, a prediction of the number counts of objects found can be seen in figure 3. We expect to detect approximately 15 such objects in our survey.




Figure 3. *Left* Number density of Ly α -emitters at redshift $z = 8.8$ to the expected flux limit of ELVIS and an EW > 30 Å. We have made 112 mock catalogs of Ly α -emitters at $z = 8.8$ by incorporating the Millennium simulation to the GALFORM semi-analytical model and extracted the number of objects detected in each catalog. This plot shows the histogram of that number. The survey volume of ELVIS is 137500 Mpc 3 , hence we expect to detect approximately 15 objects in the whole survey. *Right* Angular correlation function. Blue solid dots correspond to the median of the “measured” correlation function, red open circles correspond to the “true” correlation function, i.e. the whole catalog. Plots from Nilsson et al. (2007) in prep.

These Ly α -emitters can then be used to study the re-ionisation state of the Universe at this redshift, and to study clustering, as seen in figure 3. All of the emission lines surveyed with ELVIS are also good star formation rate indicators. Thus, the survey will enable a detailed study of the star formation history of the Universe. For H α , the survey will reach more than three orders of magnitude below L^* and so ELVIS will probe the very faint end of the H α luminosity function at $z = 0.8$.

4. Summary

To summarise, we have an extensive survey plan underway to determine the nature of Ly α -emitters at redshifts $z \sim 2 - 3$, using public, multi-wavelength data in surveys such as GOODS and COSMOS. We do this by e.g. SED fitting analysis and morphological analysis from HST data. In our GOODS-S survey,

we discovered a so-called Ly α -blob. Its properties are best explained by cold accretion onto a dark matter halo. This in turn means that the energy sources of these blobs are diverse, since previous authors have found that other blobs have been better explained by star formation (Steidel et al. 2000; Francis et al. 2001; Matsuda et al. 2004; Palunas et al. 2004) or AGN activity (Fynbo et al. 1999; Keel et al. 1999; Dey et al. 2005; Villar-Martin et al. 2005; Weidinger et al. 2005). Larger samples of this rare phenomenon are required before we can fully understand the underlying processes in these objects. Finally, we have also presented our future Public Survey (ELVIS), planned to start observing in the winter of 2008. It is expected to discover tens of Ly α -emitters at $z = 8.8$, as well as hundreds of intermediate redshift emission-line galaxies and thousands of H α -emitters at lower redshift.

Acknowledgments. K.N. gratefully acknowledges support from IDA - Instrumentcentre for Danish Astrophysics. The Dark Cosmology Centre is funded by the DNRF.

References

- Dekel, A. & Birnboim, Y., 2006, MNRAS, 368, 2
Dey, A., Bian, C., Soifer, B.T. et al. 2005, ApJ, 629, 654
Dijkstra, M., Haiman, Z. & Spaans, M. 2006, ApJ, 649, 14
Dijkstra, M., Haiman, Z. & Spaans, M. 2006, ApJ, 649, 37
Fardal, M.A., Katz, N., Gardner, J.P. et al., 2001, ApJ, 562, 605
Francis, P.J., Williger, G.M. & Collins, N.R. 2001, ApJ, 554, 1001
Fynbo, J.P.U., Møller, P. & Warren, S.J. 1999, MNRAS, 305, 849
Fynbo, J.P.U., Ledoux, C., Møller, P., Thomsen, B. & Burud, I. 2003, A&A, 407, 147
Giavalisco, M., Ferguson, H. C., Koekemoer, A. M. et al. 2004, ApJ, 600, L93
Haiman, Z., Spaans, M. & Quataert, E., 2000, ApJL 537, L5
Haiman, Z. & Rees, M.J., 2001, ApJ 556, 87
Hu, E.M. & McMahon, R.G. 1996, Nature, 382, 231
Iye, M., Ota, K., Kashikawa, N. et al. 2006, Nature, 443, 186
Keel, W.C., Cohen, S.H., Windhorst, R.A. & Waddington, I. 1999, AJ, 118, 2547
Malhotra, S., & Rhoads, J.E. 2002, ApJL, 565, L71
Matsuda, Y., Yamada, T., Hayashino, T. et al. 2004, AJ, 128, 569
Møller, P. and Warren, S.J. 1993, A&A, 270, 43
Nilsson, K.K., Fynbo, J.P.U., Møller, P., Sommer-Larsen, J., & Ledoux, C. 2006, A&A, 452, L23
Ohyama, Y., Taniguchi, Y., Kawabata, K.S. et al., 2003, ApJ, 591, L9
Palunas, P., Teplitz, H.I., Francis, P.J., Williger, G.M. & Woodgate, B.E. 2004, ApJ, 602, 545
Partridge, R.B. & Peebles, P.J.E. 1967, ApJ, 147, 868
Sommer-Larsen, J., 2005, in the proceedings of the “Island Universes: Structure and Evolution of Disk Galaxies” conference held in Terschelling, Netherlands, July 2005, ed. R de Jong (Springer Dordrecht), astro-ph/0512485
Steidel, C.C., Adelberger, K., Shapley, A.E. et al. 2000, ApJ, 532, 170
Taniguchi, Y., Shioya, Y. & Kakazu, Y., 2001, ApJL 562, L15
Venemans, B.P., Röttgering, H.J.A., Miley, G.K. et al. 2005, A&A, 431, 793
Villar-Martin, M., Sanchez, S.F. & de Breuck, C. 2005, MNRAS, 359, L5
Weidinger, M., Møller, P., Fynbo, J.P.U. & Thomsen, B. 2005, A&A, 436, 825